Ozone National Ambient Air Quality Standard Health Exceedances on August 13, 2016

Exceedance Locations and Levels

On Saturday, August 13, 2016, there were no exceedances in New Jersey of the new 8-hour average ozone NAAQS of 70 ppb that became effective in December 2015. The highest 8-hour average ozone concentration recorded in New Jersey on August 13, 2016 was 69 ppb at the Leonia station. The highest 1-hour average ozone concentration recorded was 80 ppb, also at the Leonia station, which is below the 1-hour ozone NAAQS of 120 ppb.

The number of days in 2016 on which exceedances of the new 8-hour ozone NAAQS of 70 ppb were recorded in New Jersey remains at twenty-one (21). By the 13th of August in 2015, there were a total of ten (10) days on which ozone exceedances were measured in New Jersey (based on the former 75 ppb NAAQS of 2008), and there were two (2) days by this same date in 2014.

There is a group of monitoring stations in designated counties of five (5) states, New York, Connecticut, Pennsylvania, Delaware and Maryland, that are included in New Jersey's ozone nonattainment areas. From this group of stations in the neighboring states, there were three (3) exceedances of the new 8-hour ozone NAAQS of 70 ppb recorded on Saturday, August 13, 2016 (see Table 1):

Table 1: Ozone NAAQS Exceedances at Other Monitoring Stations in New Jersey's Ozone Nonattainment Areas on August 13, 2016

STATE	STATION	Daily Maximum 8-Hr Average (ppb)
СТ	Greenwich	77
СТ	Madison-Beach Road	77
СТ	Westport	72

The highest 1-hour average ozone concentration recorded was 116 ppb at the Greenwich station in Connecticut, which is below the 1-hour ozone NAAQS of 120 ppb.

Saturday marks the 22nd day in 2016 on which an exceedance of the new 8-hour ozone NAAQS of 70 ppb was recorded in Connecticut. The number of days remains at eighteen (18) for New York, ten (10) for Pennsylvania, seven (7) for Delaware, and six (6) for Maryland.




Figure 1. Ozone Air Quality Index for August 13, 2016

Source: www.airnow.gov

For ozone terminology definitions see NJDEP Air Quality Planning's Glossary and Acronyms webpage: http://nj.gov/dep/baqp/glossary.html

Weather

Meteorological data from across the region showed temperatures reached into the mid 90°F's, while winds were from the south/southwest with a high pressure system over the Atlantic Ocean. Skies were mostly sunny over Connecticut. A low pressure surface trough was also in place along the coast of New Jersey and up through New York, which provided a mechanism to bring pollutants down to the surface. This weather feature, in combination with adequate sunlight, southwest winds, and warm temperatures, are all meteorological conditions commonly seen on high ozone days.

Where Did the Air Pollution that Caused Ozone Come From?

Figures 2, 3, and 4 show the back trajectories at different wind heights for the monitored exceedances on August 13, 2016. The figures illustrate where the winds came from during the 48 hours preceding the high ozone event.

The low level wind (Figure 2) traveled up the I-95 corridor and through the major metropolitan areas of Baltimore, Philadelphia, and New York City where it picked up emissions from cars, trucks, and industry on the way to the exceedance monitors in Connecticut. The back trajectories for the mid-level (Figure 3) and higher level (Figure 4) winds illustrate similar transport pathways. Winds originated over the Smoky Mountains and traveled across Pennsylvania, northern New Jersey, and New York City, bringing

additional emissions from motor vehicles, industry, and power plants to Connecticut. These winds, in combination with the low level wind, caused air pollution from a variety of mobile and stationary sources to be transported in the areas of coastal Connecticut that experienced high ozone on August 13, 2016.

Figure 2. 48-hour Back Trajectories for August 13, 2016 at 10 meters

NOAA HYSPLIT MODEL Backward trajectories ending at 1800 UTC 13 Aug 16 NAM Meteorological Data




Figure 3. 48-hour Back Trajectories for August 13, 2016 at 500 meters

NOAA HYSPLIT MODEL Backward trajectories ending at 1800 UTC 13 Aug 16 NAM Meteorological Data





Figure 4. 48-hour Back Trajectories for August 13, 2016 at 1500 meters

NOAA HYSPLIT MODEL Backward trajectories ending at 1800 UTC 13 Aug 16 NAM Meteorological Data



How is Smog Created?

Ground-level ozone, also known as smog, is an air pollutant known to cause a number of health effects and negatively impact air quality and the environment in the state of New Jersey. Smog is formed when oxides of nitrogen (NOx) and volatile organic compounds (VOCs) react in the presence of sunlight. Smog can irritate any set of lungs, but those with lung-related deficiencies should take extra precautions on bad ozone days.

Find Out About Air Quality Every Day

The "What's Your Air Quality Today?" page at http://www.nj.gov/dep/cleanairnj/ tells you how to sign up to receive notifications and find out when your local air has reached unhealthy ozone levels.