
Data Summary of the Delaware River Seine Survey 2023





In 2023, biologists conducted the survey from June 19 to October 26, 2023. During those five months, crews hauled 287 individual seines catching 34,170 total fish, averaging 119 fish per haul. The five most abundant species caught were: bay anchovy, banded killifish, Atlantic menhaden, spottail shiner and Eastern silvery minnow.

In 2023, striped bass was the fourteenth most abundant fish caught. In total, 325 striped bass were caught and 232 of those were young-of-year (YOY) striped bass meaning that 71% of the striped bass caught were less than 1 year old. The YOY index (geometric mean) for 2023 was 0.26, which ranks thirty-eighth highest out of the 44-year time series.



All water quality parameters for 2023 were within normal range. Water temperatures ranged from 13.1 to 30.4 degrees Celsius. Dissolved oxygen levels ranged from 0.2 to 9.8 milligrams per liter. Salinity values ranged from 0 to 12.5 parts per thousand. pH values collected averaged 7.6, which is considered slightly more basic than neutral (7.0).

Catch totals by species for 2023 can be found below:

2023 CATCH TOTALS

| COMMON NAME | SCIENTIFIC NAME | TOTAL NUMBER | FREQUENCY OF OCCURRENCE (%) |
|------------------------|-------------------------------------|--------------|-----------------------------|
| Alewife | Alosa pseudoharengus | 18 | 2.439% |
| American Eel | Anguilla rostrata | 24 | 7.317% |
| American Shad | Alosa sapidissima | 1,092 | 39.721% |
| Atlantic Croaker | Micropogonias undulatus | 445 | 16.725% |
| Atlantic Menhaden | Brevoortia tyrannus | 3,384 | 22.648% |
| Atlantic Needlefish | Strongylura marina | 5 | 1.394% |
| Atlantic Silverside | Menidia menidia | 379 | 20.557% |
| Banded Killifish | Fundulus diaphanus | 6,071 | 54.007% |
| Banded Sunfish | Enneacanthus obesus | 1 | 0.348% |
| Bay Anchovy | Anchoa mitchilli | 10,904 | 40.418% |
| Black Bullhead | Ameiurus melas | 1 | 0.348% |
| Black Drum | Pogonias cromis | 6 | 1.394% |
| Blackcheek Tonguefish | Symphurus plagiusa | 1 | 0.348% |
| Blue Crab | Callinectes sapidus | 301 | 25.087% |
| Blueback Herring | Alosa aestivalis | 659 | 13.240% |
| Bluefish | Pomatomus saltatrix | 33 | 8.362% |
| Bluegill Sunfish | Lepomis macrochirus | 52 | 8.711% |
| Brown Bullhead | Ameiurus nebulosus | 1 | 0.348% |
| | | 8 | 1.742% |
| Carp Channel Catfish | Cyprinus carpio Ictalurus punctatus | 36 | 5.923% |
| | · | | |
| Comely Shiner | Notropis amoenus | 6 | 0.697% |
| Crevalle Jack | Caranx hippos | 4 | 1.045% |
| Eastern Silvery Minnow | Hybognathus regius | 2,476 | 35.540% |
| Flathead Catfish | Pylodictis olivaris | 1 | 0.348% |
| Gizzard Shad | Dorosoma cepedianum | 239 | 19.861% |
| Golden Shiner | Notemigonus crysoleucas | 3 | 1.045% |
| Goldfish | Carassius auratus | 7 | 2.091% |
| Green Sunfish | Lepomis cyanellus | 1 | 0.348% |
| Hogchoker | Trinectes maculatus | 515 | 22.648% |
| Inland Silverside | Menidia beryllina | 239 | 16.028% |
| Largemouth Bass | Micropterus salmoides | 9 | 2.091% |
| Mummichog | Fundulus heteroclitus | 1,344 | 29.965% |
| Northern Hog Sucker | Hypentelium nigricans | 23 | 4.181% |
| Northern Snakehead | Channa argus | 1 | 0.348% |
| Pumpkinseed Sunfish | Lepomis gibbosus | 20 | 3.484% |
| Rainwater Killifish | Lucania parva | 1 | 0.348% |
| Redbreast Sunfish | Lepomis auritus | 4 | 0.697% |
| Rough Silverside | Membras martinica | 179 | 9.059% |
| Smallmouth Bass | Micropterus dolomieu | 28 | 6.969% |
| Spanish Mackerel | Scomberomorus maculatus | 2 | 0.697% |
| Spot | Leiostomus xanthurus | 136 | 13.240% |
| Spottail Shiner | Notropis hudsonius | 3,318 | 38.328% |
| Striped Anchovy | Anchoa hepsetus | 21 | 3.484% |
| Striped Bass | Morone saxatilis | 325 | 31.010% |
| Striped Killifish | Fundulus majalis | 2 | 0.697% |
| Summer Flounder | Paralichthys dentatus | 4 | 1.045% |
| Tessellated Darter | Etheostoma olmstedi | 769 | 33.101% |
| Walleye | Sander vitreus | 2 | 0.348% |
| Weakfish | Cynoscion regalis | 6 | 0.348% |
| White Catfish | Ameiurus catus | 3 | 1.045% |
| White Perch | Morone americana | 987 | 49.826% |
| White Sucker | Catostomus commersonii | 60 | 6.272% |
| Yellow Bullhead | Ameiurus natalis | 6 | 0.348% |
| Yellow Perch | Perca flavescens | 8 | 1.394% |
| GRAND TOTAL | | 34,170 | |