Parent Threat	Level 2	Priority Level 3 Threats	Inner Coastal Plain Level 3 Priority Threats
	2 Agriculture and Aquaculture: Threats from agricultural activities, such as the expansion and intensification of agriculture and livestock farming, including silviculture, mariculture and aquaculture and related infrastructures. This includes the initial conversion of habitat (deforestation, filling/excavation, draining of wetlands, etc.) that is associated with cultivation or infrastructure development, as well as uses and practices (intensification of agricultural practices, use of machinery, etc.), but not the transport of the resources that are produced, crop irrigation or pollution.		
	2.1	vineyards, mix	erennial Non-Timber Crops: Non-timber crops that are planted for food, fodder, fuel or other uses; farms, crop fields, ed agroforestry systems, etc. For rotational crops, consider the most intensive practice that is used. Considering the diversity of actices and related impacts, some speciality cultures will be pooled into a generic threat category.
		2.1.1	Annual cropping systems (field crops): Wide-row crops that require the most intensive agricultural practices and which have the most significant impacts. E.g., maize (corn), soybean, barley, vegetable crops, oats, wheat, canola, hemp.
	7 Natural Systems Modifications: Threats from activities that are generally carried out to improve human welfare, but may result in habitat degradation or destruction. This threat category includes the development or redevelopment (management) of natural and semi-natural habitats, as well as certain natural processes that can act as threats. Stopping a conservation action or a practice that is conducive to conservation is not interpreted as a threat, but as a return to the source threat (e.g., vegetation succession affecting pioneer species). Excludes meteorological or climate change-related threats that may modify natural systems.		
	7.2	Dams and Wa	ter Management/Use: Facilities or activities that alter the natural water regime (flow or water levels).
		7.2.3	Water management using culverts: The design, installation and management of culverts that are used to permit water flow under roads or railroads can cause discontinuities in streams and promote erosion.
		7.2.7	Withdrawal of groundwater: Withdrawal of groundwater for human consumption, crop production or other purposes. E.g., pumping water from the water table.
	7.3		em Modifications: Other activities that contribute to habitat alteration or loss by redeveloping natural systems to improve human distinguished from the development and maintenance of urban parks.
		7.3.2	Vegetation succession: Natural vegetation succession causing habitat loss for species of early successional habitats.
	7.4		educing Human Maintenance: Stopping, reducing, or removing a management activity. Includes vegetation control, FIRE , hydrology control, human disturbance, etc.
8		7.4.1	Reducing or ceasing vegetation control: e.g. reducing, removing, or ceasing prescribed fire, removal of invasive species, maintenance of early successional vegetation
	materials) that	have or are exp Invasive Non-	tic Species, Genes and Diseases: Threats posed by non-native and native species (plants, animals, pathogens or genetic ected to have harmful effects on biodiversity following their introduction, spread or increase in population (abundance). Native / Alien Plants & Animals: Harmful plants and animals that were not originally present within an ecosystem, but were rectly introduced into or spread in the ecosystem as a result of human activities. The concept of exotic species includes species
		that are not na US. Domestic s	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also
		that are not na US. Domestic s includes introd	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases".
		that are not na US. Domestic s includes introd 8.1.1	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also fuction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling
		that are not na US. Domestic s includes introd 8.1.1 8.1.2	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases".
	8.2	that are not na US. Domestic s includes introd 8.1.1 8.1.2 8.1.4 Problematic N	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife.
		that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese.
		that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biole	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations
		that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biole	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the species are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese.
9	8.5 Pollution: Three pollution are ty	that are not na US. Domestic s includes introd 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 reats that are ass pically correlat ough there is a c	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also function of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. Aquatic plants: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks . ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by
9	Pollution: Three pollution are ty effluents). Although the source activity of the source ac	that are not na US. Domestic s includes introd 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 eats that are ass pically correlat ugh there is a d vity.	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also function of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. Aquatic plants: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks . ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural
9	Pollution: Three pollution are ty effluents). Although the source activity of the source ac	that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 Easts that are ass pically correlat ugh there is a c vity. Domestic and nutrients, sedin	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the species are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also huttor of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. Intereased grazing by the chestnut, European water milfoil, hydrilla, purple loosestrife. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. Increased grazing by vertebrates: E.g., population isolation, inbreeding, bottlenecks . ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural lirect correlation between pollution and these other threats, their impact (scope and severity) is often evaluated separately from Urban Waste Water: Point or non-point source wastewater from residential and urban areas; these discharges (may) contain ments, toxic substances, chemicals, etc.
9	Pollution: Three pollution are ty effluents). Although the source activity of the source ac	that are not na US. Domestic s includes introd 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 Turtinsic Biolo 8.5.1 stats that are ass pically correlat ough there is a c vity. Domestic and nutrients, sedin 9.1.1	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the species are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also fuction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks . ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural lirect correlation between pollution and these other threats, their impact (scope and severity) is often evaluated separately from nents, toxic substances, chemicals, etc. Domestic wastewater: Liquid domestic waste that is produced by urban centres and discharged primarily by the sewage system. E.g., discharges from municipal waste treatment plants, leaks from sewars/septic tanks, untreated discharges, pit toilets, medical
9	8.5 Pollution: Three pollution are typ effluents). Altho the source activ 9.1	that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 ats that are ass pically correlat pugh there is a d vity. Domestic and nutrients, sedir 9.1.1 9.1.2 Agricultural a discharges are	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the species are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also neution of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased e they are new "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks . ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural lirect correlation between pollution and these other threats, their impact (scope and severity) is often evaluated separately from untan activities listed in the other sections (e.g., air pollution areas; these discharges (may) contain ments, toxic substances, chemicals, etc. Domestic wastewater: Liquid domestic waste that is produced by urban centres and discharged primarily by the sewage system. E.g., discharges from municipal waste treatment plants, leaks from sewers/septic tanks, untreated discharges, pit toilets, medical components in water (birth control hormones, antidepressants, antibiotics), toxoplasmosis, etc. Run-off: Effluents resulting from urban activities that are separate from the water supply system. Excludes oils and other
9	8.5 Pollution: Three pollution are typ effluents). Altho the source activ 9.1	that are not na US. Domestic s includes introd 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 ats that are ass pically correlat pugh there is a d vity. Domestic and nutrients, sedir 9.1.1 9.1.2 Agricultural a discharges are chemicals, etc. machinery.	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. ative Plants & Animals: Plants and animals that were originally present in ecosystem(s), but whose populations have increased et they are now "out of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks. ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural lirect correlation between pollution and these other threats, their impact (scope and severity) is often evaluated separately from Urban Waste Water: Point or non-point source wastewater from residential and urban areas; these discharges (may) contain nents, toxic substances, chemicals, etc. Domestic wastewater: Liquid domestic waste that is produced by urban centres and discharged primarily by the sewage system E.g., discharges from municipal waste treatment plants, leaks from sewers/septic tanks, untreated discharges, pit toilets, medical components in water (birth control hormones, antidepressants, antibiotics), toxoplasmosis, etc. Run-off: Effluents resulting from urban activities that are separate from th
9	8.5 Pollution: Three pollution are typ effluents). Altho the source activ 9.1	that are not na US. Domestic s includes introa 8.1.1 8.1.2 8.1.4 Problematic N to a level wher 8.2.2 Intrinsic Biolo 8.5.1 ats that are ass pically correlat pugh there is a d vity. Domestic and nutrients, sedir 9.1.1 9.1.2 Agricultural a discharges are chemicals, etc. machinery. 9.3.1	tive to a specific habitat; it can therefore include the introduction of species that are considered native to a different region of the pecies are also considered nonnative, whether they are feral or semi-domesticated (e.g., domestic cats going outside). Also uction of wildlife due to "mercy releases". Terrestrial animals: E.g., Norway rats, feral cats, European Starling Terrestrial plants: E.g., Norway rats, feral cats, European Starling Aquatic plants: E.g., Norway rats, feral cats, European Starling Aquatic plants: E.g., Rosa multiflora, garlic mustard, Japanese stiltgrass, Phragmites. Aquatic plants: E.g., Water chestnut, European water milfoil, hydrilla, purple loosestrife. Introduction of control" or overabundant as a direct or indirect result of certain human activities. Increased grazing by vertebrates: E.g., increased grazing by white-tailed deer and snow geese. gical Limitations Loss of Genetic Diversity: e.g. population isolation, inbreeding, bottlenecks. ociated with the introduction of foreign or excess material/energy from point and non-point sources. Threats that are posed by ed with other human activities listed in the other sections (e.g., air pollution from cars, water pollution from sewage, agricultural litect correlation between pollution and these other threats, their impact (scope and severity) is offen evaluated separately from Urban Waste Water: Point or non-point source wastewater from residential and urban areas; these discharges (may) contain nents, toxic substances, chemicals, etc. Domestic wastewater: Liquid domestic waste that is produced by urban centres and discharged primarily by the sewage system E.g., discharges from municipal waste treatment plants, leaks from sewers/septic tanks, untreated discharges, pit toilets, medical components in water (birth control hormones, antidepressants, antibiotics), toxoplasmosis, etc. Run-off: Effluents: Wastewater (pollutants) that is generated by a